

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number  
**WO 01/96338 A1**(51) International Patent Classification<sup>7</sup>: **C07D 471/14**,  
A61K 31/55 // (C07D 471/14, 243:00, 221:00)[CA/CA]; 2615, de la Volière, Laval, Québec H7L 3V6  
(CA).

(21) International Application Number: PCT/CA01/00890

(74) Agent: **BERNIER, Louise, G.**; Boehringer Ingelheim  
(Canada) Ltd., 2100 Cunard, Laval, Québec H7S 2G5  
(CA).

(22) International Filing Date: 14 June 2001 (14.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/212,329 16 June 2000 (16.06.2000) US  
60/256,638 18 December 2000 (18.12.2000) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(71) Applicant (*for all designated States except US*):  
**BOEHRINGER INGELHEIM (CANADA) LTD.**  
[CA/CA]; 2100 Cunard Street, Laval, Québec H7S 2G5  
(CA).(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventor; and


Published:

(75) Inventor/Applicant (*for US only*): **SIMONEAU, Bruno**





— with international search report

[Continued on next page]

(54) Title: NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS



(57) Abstract: Provided are compounds of general formula (I), wherein R<sup>2</sup> is selected from the group consisting of H, F, Cl, (C<sub>1-4</sub>) alkyl, (C<sub>3-4</sub>) cycloalkyl and CF<sub>3</sub>; R<sup>4</sup> is H or Me; R<sup>5</sup> is H, Me or Et, with the proviso that R<sup>4</sup> and R<sup>5</sup> are not both Me, and if R<sup>4</sup> is Me then R<sup>5</sup> cannot be Et; R<sup>11</sup> is Et, cyclopropyl, propyl, isopropyl, or isobutyl; and Q is selected from the group consisting of (II), (III), (IV) and (V); and pharmaceutically acceptable salts thereof, as inhibitors of HIV reverse transcriptase, wild-type and several mutant strains.



WO 01/96338 A1



— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS****TECHNICAL FIELD OF THE INVENTION**

5

The invention relates to novel compounds and pharmaceutically acceptable salts thereof, their use, either alone or in combination with other therapeutic agents, in the treatment or prophylaxis of HIV infection, and to pharmaceutical compositions comprising the compounds.

10

**BACKGROUND OF THE INVENTION**

The disease known as acquired immune deficiency syndrome (AIDS) is caused by the human immunodeficiency virus (HIV), particularly the strain known as HIV-1. In order for HIV to be replicated by a host cell, the information of the viral genome must be integrated into the host cell's DNA. However, HIV is a retrovirus, meaning that its genetic information is in the form of RNA. The HIV replication cycle therefore requires a step of transcription of the viral genome (RNA) into DNA, which is the reverse of the normal chain of events. The transcription of the viral RNA into DNA is accomplished by an enzyme that has been aptly dubbed reverse transcriptase (RT). The HIV virion includes a copy of RT along with the viral RNA.

Reverse transcriptase has three known enzymatic functions; it acts as an RNA-dependent DNA polymerase, as a ribonuclease, and as a DNA-dependent DNA polymerase. Acting as an RNA-dependent DNA polymerase, RT transcribes a single-stranded DNA copy of the viral RNA. Acting as a ribonuclease, RT destroys the original viral RNA, and frees the DNA just produced from the original RNA. Finally, acting as a DNA-dependent DNA polymerase, RT makes a second, complementary DNA strand, using the first DNA strand as a template. The two strands form

30

double-stranded DNA, which is integrated into the host cell's genome by another enzyme called integrase.

Compounds that inhibit the enzymatic functions of HIV-1 reverse transcriptase will inhibit replication of HIV-1 in infected cells. Such compounds are useful in the prevention or treatment of HIV-1 infection in human subjects, as demonstrated by known RT inhibitors such as 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxyinosine (ddI), 2',3'-dideoxycytidine (ddC), d4T, 3TC, Nevirapine, Delavirdine, Efavirenz and Abacavir, the main drugs thus far approved for use in the treatment of AIDS.

As with any antiviral therapy, use of RT inhibitors in the treatment of AIDS eventually leads to a virus that is less sensitive to the given drug. Resistance (reduced sensitivity) to these drugs is the result of mutations that occur in the reverse transcriptase segment of the *pol* gene. Several mutant strains of HIV have been characterised, and resistance to known therapeutic agents is due to mutations in the RT gene. Some of the most commonly observed mutants clinically are: the Y181C mutant, in which a tyrosine (Y), at codon 181, has been mutated to a cysteine (C) residue, and K103N where the lysine (K) at position 103 has been replaced by asparagine (N). Other mutants which emerge with increasing frequency during treatment with known antivirals include the single mutants V106A, G190A, Y188C, and P236L; and the double mutants K103N/Y181C, K103N/P225H, K103N/V108I, and K103N/L100I.

As therapy and prevention of HIV infection using antivirals continues, the emergence of new resistant strains is expected to increase. There is therefore an ongoing need for new inhibitors of RT, with different patterns of effectiveness against the various mutants.

Compounds having tricyclic structures which are inhibitors of HIV-1 are

described in U.S. Pat. No. 5,366,972. Other inhibitors of HIV-1 reverse transcriptase are described in Hargrave et al., J. Med Chem., 34, 2231 (1991).


U.S. Pat. No. 5,705,499 proposes 8-arylalkyl- and 8-arylheteroalkyl-5,11-dihydro-6H-dipyrido[3,2-B:2',3'-E][1,4]diazepines as inhibitors of RT. The exemplified compounds are shown to have some activity against wild type and mutant HIV-1 RT, particularly Y181C and other single mutants such as K103N albeit less effectively.

Specifically, the compounds of the present invention are effective at inhibiting the Y181C and K103N mutants as well as a broad range of other single mutants and certain commonly found double mutants such as K103N/Y181C and K103N/P225H.

#### SUMMARY OF THE INVENTION

The invention reduces the difficulties and disadvantages of the prior art by providing novel compounds that are potent inhibitors of single and double mutant strains of HIV-1 RT.

In a first aspect the invention provides a compound of the general formula I:



wherein


R<sup>2</sup> is selected from the group consisting of H, F, Cl, C<sub>1-4</sub> alkyl, C<sub>3-4</sub> cycloalkyl and CF<sub>3</sub>;

R<sup>4</sup> is H or Me;

R<sup>5</sup> is H, Me or Et, with the proviso that R<sup>4</sup> and R<sup>5</sup> are not both Me, and if R<sup>4</sup> is Me then R<sup>5</sup> cannot be Et;

R<sup>11</sup> is Me, Et, cyclopropyl, propyl, isopropyl, or cyclobutyl;

5 Q is selected from the group consisting of:



or a pharmaceutically acceptable salt thereof.

10 In a second aspect, the invention provides an inhibitor of HIV replication, of the general formula I, or a pharmaceutically acceptable salt thereof.

In a third aspect, the invention provides an inhibitor of a reverse transcriptase enzyme of HIV, of the general formula I, or a pharmaceutically acceptable salt thereof.

15

In a fourth aspect, the invention provides a method for the treatment or prevention of HIV infection, comprising administering to a patient an HIV inhibiting amount of a compound of formula I, or a pharmaceutically acceptable salt thereof.

20

In a fifth aspect, the invention provides a pharmaceutical composition for the

treatment or prevention of HIV infection, comprising a compound of formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

- 5 In a sixth aspect, the invention provides a method for preparation of a compound of formula I, or a pharmaceutically acceptable salt thereof.

#### **DETAILED DESCRIPTION OF THE INVENTION**

##### **Definitions**

- 10 As used herein, the term "C<sub>1-4</sub> alkyl" is intended to mean linear or branched alkyl radicals containing from one to four carbon atoms and includes methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl and *tert*-butyl.

- As used herein, the term "C<sub>3-4</sub> cycloalkyl" is intended to mean saturated  
15 cyclic hydrocarbon radicals containing three to four carbon atoms and includes cyclopropyl and cyclobutyl.

##### **Detailed description of preferred embodiments**

- According to a preferred embodiment, compounds of the invention are defined according to formula I wherein R<sup>2</sup> is preferably Cl, F, or H. More  
20 preferably, R<sup>2</sup> is Cl or H. Most preferably, R<sup>2</sup> is H.

According to a preferred embodiment, compounds of the invention are defined according to formula I wherein R<sup>4</sup> is preferably H.


- 25 According to an alternative embodiment, compounds of the invention are defined according to formula I wherein preferably R<sup>5</sup> is Me.

Preferably, compounds of the invention are defined according to formula I wherein R<sup>11</sup> is Et or cyclopropyl. More preferably, R<sup>11</sup> is Et.




6


According to a preferred embodiment, compounds of the invention are defined according to formula I wherein Q is preferably selected from the group consisting of:




More preferably, Q is:



10 Alternatively, preferred embodiments of the invention include compounds selected from the group consisting of:



7



The compounds of the invention are effective inhibitors of wild type reverse transcriptase as well as inhibiting, for example, the single mutation enzymes Y181C, K103N, V106A, G190A, Y188C, and P236L. The compounds also  
5 inhibit the double mutation enzymes K103N/Y181C, K103N/P225H, K103N/V108I and K103N/L100I.

The compounds of formula I possess inhibitory activity against HIV-1 reverse transcriptase. When administered in suitable dosage forms, they are  
10 useful in the treatment of AIDS, ARC and related disorders associated with HIV-1 infection. Another aspect of the invention, therefore, is a method for treating HIV-1 infection which comprises administering to a human being, infected by HIV-1, a therapeutically effective amount of a novel compound of formula I, as described above. Whether it be termed treatment or  
15 prophylaxis, the compounds may also be used to prevent perinatal transmission of HIV-1 from mother to baby, by administration to the mother prior to birth.

The compounds of formula I may be administered in single or divided doses  
20 by the oral or parenteral routes. A suitable oral dosage for a compound of formula I would be in the range of about 0.5 mg to 1 g per day. A preferred oral dosage for a compound of formula I would be in the range of about 100 mg to 800 mg per day for a patient weighing 70 kg. In parenteral formulations, a suitable dosage unit may contain from 0.1 to 250 mg of said  
25 compounds, preferably 1 mg to 200 mg. It should be understood, however,

that the dosage administration from patient to patient will vary and the dosage for any particular patient will depend upon the clinician's judgement, who will use as criteria for fixing a proper dosage the size and condition of the patient as well as the patient's response to the drug.

5

When the compounds of the present invention are to be administered by the oral route, they may be administered as medicaments in the form of pharmaceutical preparations, which contain them in association with a compatible pharmaceutical carrier material. Such carrier material can be an inert organic or inorganic carrier material suitable for oral administration. Examples of such carrier materials are water, gelatin, talc, starch, magnesium stearate, gum arabic, vegetable oils, polyalkylene-glycols, petroleum jelly and the like.

15 The pharmaceutical preparations can be prepared in a conventional manner and finished dosage forms can be solid dosage forms, for example, tablets, dragees, capsules, and the like, or liquid dosage forms, for example solutions, suspensions, emulsions and the like. The pharmaceutical preparations may be subjected to conventional pharmaceutical operations such as sterilisation.

20 Further, the pharmaceutical preparations may contain conventional adjuvants such as preservatives, stabilisers, emulsifiers, flavour-improvers, wetting agents, buffers, salts for varying the osmotic pressure and the like. Solid carrier material which can be used include, for example, starch, lactose, mannitol, methyl cellulose, microcrystalline cellulose, talc, silica, dibasic

25 calcium phosphate, and high molecular weight polymers (such as polyethylene-glycol).

For parenteral use, a compound of formula I can be administered in an aqueous or non-aqueous solution, suspension or emulsion in a

pharmaceutically acceptable oil or a mixture of liquids, which may contain bacteriostatic agents, antioxidants, preservatives, buffers or other solutes to render the solution isotonic with the blood, thickening agents, suspending agents or other pharmaceutically acceptable additives. Additives of this type  
5 include, for example, tartrate, citrate and acetate buffers, ethanol, propylene glycol, polyethylene glycol, complex formers (such as EDTA), antioxidants (such as sodium bisulfite, sodium metabisulfite, and ascorbic acid), high molecular weight polymers (such as liquid polyethylene oxides) for viscosity regulation and polyethylene derivatives of sorbitol anhydrides. Preservatives  
10 may also be added if necessary, such as benzoic acid, methyl or propyl paraben, benzalkonium chloride and other quaternary ammonium compounds.


The compounds of this invention may also be administered as solutions for  
15 nasal application and may contain in addition to the compounds of this invention suitable buffers, tonicity adjusters, microbial preservatives, antioxidants and viscosity-increasing agents in an aqueous vehicle. Examples of agents used to increase viscosity are polyvinyl alcohol, cellulose derivatives, polyvinylpyrrolidone, polysorbates or glycerin. Microbial  
20 preservatives added may include benzalkonium chloride, thimerosal, chlorobutanol or phenylethyl alcohol.

Additionally, the compounds provided by the invention can be administered by suppository.


25

The compounds of the invention may be made using the skills of a synthetic organic chemist. An exemplary reaction scheme is shown in schemes 1 to 6.

**Scheme 1: Preparation of intermediates in which R<sup>4</sup> is Me**





**Scheme 2: Preparation of intermediates in which R<sup>5</sup> is Me**





- 5 The sequence of scheme 2 is analogous to one described by J.M. Klunder *et al.*; *J. Med. Chem.* **1998**, 41, 2960-71, and C.L. Cywin *et al.*; *J. Med. Chem.* **1998**, 41, 2972-84.

11

**Scheme 3: Preparation of intermediates in which R<sup>2</sup> is C<sub>1-3</sub> alkyl****Scheme 4: Preparation of intermediates in which R<sup>2</sup> is CF<sub>3</sub>**

## 12

**Scheme 5: Introduction of the quinoline nucleus****Scheme 6: Alternate route to compounds in which R<sup>4</sup> is Me**

5

As stated before, the compounds provided by the invention inhibit the enzymatic activity of HIV-1 RT. Based upon testing of these compounds, as described below, it is known that they inhibit the RNA-dependent DNA polymerase activity of HIV-1 RT. It is known (data not shown) that they also inhibit the DNA-dependent DNA polymerase activity of HIV-1 RT. Utilising the Reverse Transcriptase (RT) Assay described below, compounds can be

tested for their ability to inhibit the RNA-dependent DNA polymerase activity of HIV-1 RT. Certain specific compounds described in the Examples which appear below, were so tested. The results of this testing appear in Table 2 as IC<sub>50</sub> (nM) and Table 3 as EC<sub>50</sub> (nM).

5

### EXAMPLES

The present invention is illustrated in further detail by the following non-limiting examples. All reactions were performed in a nitrogen or argon  
10 atmosphere. Temperatures are given in degrees Celsius. Solution percentages or ratios express a volume to volume relationship, unless stated otherwise.

Abbreviations or symbols used herein include:

- DEAD: diethyl azodicarboxylate;
- 15 DIAD: diisopropyl azodicarboxylate;
- DIEA: diisopropylethylamine;
- DMAP: 4-(dimethylamino)pyridine;
- DMSO: dimethylsulfoxide;
- DMF: dimethylformamide;
- 20 ES MS: electron spray mass spectrometry;
- Et: ethyl;
- EtOAc: ethyl acetate;
- Et<sub>2</sub>O: diethyl ether;
- HPLC: high performance liquid chromatography;
- 25 iPr: isopropyl;
- Me: methyl;
- MeOH: methanol;
- MeCN: acetonitrile;
- NBS: N-bromosuccinimide;



- Ph: phenyl;  
TBE: tris-borate-EDTA;  
TBTU: 2-(1*H*-benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium  
tetrafluoroborate;  
5 TFA: trifluoroacetic acid;  
THF: tetrahydrofuran;  
MS (ES): electrospray mass spectrometry;  
MS (FAB) or FAB/MS: fast atom bombardment mass spectrometry;  
HRMS: high resolution mass spectrometry;  
10 PFU: plaque forming units;  
DEPC: diethyl pyrocarbonate;  
DTT: dithiothreitol;  
EDTA: ethylenediaminetetraacetate;  
UMP: uridine 5'-monophosphate;  
15 UTP: uridine 5'-triphosphate;  
MES: 2-(*n*-morpholino)ethanesulfonic acid;  
SDS-PAGE: sodium dodecyl sulfate-polyacrylamide gel electrophoresis;  
MWCO: molecular weight cut-off;  
Bis-Tris Propane: 1,3-Bis{tris(hydroxymethyl)-methylamino}propane;  
20 GSH: reduced glutathione;  
OBG: *n*-Octyl- $\beta$ -D-glucoside.

### SYNTHESSES

- The following examples illustrate methods for preparing compounds of the  
25 invention.

#### Example I (entry 24, Table 1)

*5,11-Dihydro-11-ethyl-5-methyl-8-[2-(4-quinolinylloxy)ethyl]-6H-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one*

**a) 2-(Ethylamino)-3-nitropyridine**

To a solution of 2-chloro-3-nitropyridine (51 g, 325 mmol) in THF (650 mL) was added a 2 M solution of ethylamine in THF (365 mL, 731 mmol). The reaction was stirred at room temperature overnight. The reaction mixture was poured into water (~1.5 L) and the resulting solid was filtered and dried under reduced pressure to give the title compound (52 g).

**b) 3-Amino-2-(ethylamino)pyridine**

A solution of 2-(ethylamino)-3-nitropyridine (52 g) in MeOH (600 mL) was stirred overnight at room temperature under hydrogen (1 atm.) in the presence of 20% Pd(OH)<sub>2</sub>/C (10.4 g). The catalyst was removed by filtration through diatomaceous earth. The filtrate was concentrated under reduced pressure to give the title compound as a black solid (39 g, 88% yield over steps a) and b).

**c) 2-Chloro-N-[2-(ethylamino)-3-pyridinyl]-5-bromo-3-pyridinecarboxamide**

To a cooled solution of 3-amino-2-(ethylamino)pyridine (30.6 g, 223 mmol) in MeCN (740 mL) was added solid NaHCO<sub>3</sub> (56.3 g, 669 mmol). After 5 min, crude 5-bromo-2-chloro-3-pyridinecarbonyl chloride (prepared from 5-bromo-2-hydroxy-3-pyridinecarboxylic acid and SOCl<sub>2</sub> [as described by T. W. Gero *et al.* in *Synth. Commun.* 1989; 19, 553-559 (incorporated herein by reference) but with omission of the aqueous work-up] was added (1 equiv., 223 mmol). After 2 h, the reaction mixture was poured over ice/H<sub>2</sub>O (1.5 L) and the resulting solid was filtered, rinsed with H<sub>2</sub>O and then hexane. After drying under reduced pressure overnight, the title compound was obtained as a black solid (54.9 g, 69% yield).

**d) 8-Bromo-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

To a solution of 2-chloro-N-[2-(ethylamino)-3-pyridinyl]-5-bromo-3-

pyridinecarboxamide (54.9 g, 154.4 mmol) in pyridine (308 mL) at 50 °C was added drop-wise a 1 M solution of NaHMDS (sodium hexamethyldisilazide) in THF (355 mL, 355 mmol). After 10 min, the reaction was allowed to cool to room temperature, and then was poured over ice water (2 L). The resulting solid was filtered, rinsed with water and then hexane. The solid was dried under reduced pressure to give the title compound (36 g, 75% yield) as a dark green solid.

**e) 8-Bromo-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

To a solution of the 8-bromo-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (36.7 g, 115 mmol) in DMF (380 mL) was added NaH (3.5 g, 138 mmol), and the mixture was heated to 50 °C for 30 min. The reaction mixture was cooled to room temperature and treated with MeI (14.3 mL, 230 mmol). After 1.5 h, the reaction mixture was poured over ice water. The solid was filtered, washed with water and then hexane to give after drying the title compound (37.9 g 99% yield) as a dark grey solid.

**f) 5,11-Dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

Allyltributyltin (30.7 mL, 99.0 mmol) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (5.20 g, 4.50 mmol) were added to a degassed (N<sub>2</sub> through solution for 30 min) solution of 8-bromo-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (30.0 g, 90.0 mmol) in DMF (450 mL) at room temperature. The mixture was stirred at 90 °C for 1.5 h then was cooled to room temperature and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane: EtOAc, 8:2 to 7:3) to give the title compound (22.19 g, 84% yield).

**g) 5,11-Dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-**

**b:2',3'-e][1,4]diazepin-6-one**

A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of 5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6*H*-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (22.19 g, 75.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (150 mL) and MeOH (150 mL) for 2.5 h. A stream of N<sub>2</sub> was next bubbled through the solution for 15 min and then solid NaBH<sub>4</sub> (4.99 g, 132 mmol) was added to the solution. The reaction mixture was allowed to warm to room temperature. After 1 h, aqueous saturated NH<sub>4</sub>Cl (200 mL) was added and the mixture was stirred at room temperature for 2 h. The organic solvents were removed under reduced pressure. Water (300 mL) and CHCl<sub>3</sub> (300 mL) were added to the residue. The phases were separated and the aqueous layer was extracted with CHCl<sub>3</sub> (3 × 300 mL). The combined organic layers were dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:CHCl<sub>3</sub>, 4:1) to give the title compound (16.1 g, 72% yield) as a white solid.

**h) 5,11-Dihydro-11-ethyl-5-methyl-8-[2-(4-quinolinylloxy)ethyl]-6*H*-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

Diethyl azodicarboxylate (DEAD) (12.8 mL, 81.0 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6*H*-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (16.1 g, 54.0 mmol), 4-hydroxyquinoline (11.6 g, 81.0 mmol) and Ph<sub>3</sub>P (21.3 g, 81.0 mmol) in THF (270 mL) at room temperature. The mixture was stirred at room temperature for 1h then was concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH; 95:5) to give the title compound (17.7 g, 77% yield) as a white solid: MS (ESI) *m/z* 426 (MH)<sup>+</sup>.

**Example II (entry 2, Table 1)**

**2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-[2-(4-quinolinylloxy)ethyl]-6*H*-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

**a) 6-Chloro-2-(ethylamino)-3-nitropyridine**

An ice-cold solution of EtNH<sub>2</sub> (49.8 g, 1.10 mol) in toluene (200 mL) was added over 15 min to an ice-cold solution of 2,6-dichloro-3-nitropyridine (100.0 g, 0.52 mol) in toluene (225 mL). The mixture was stirred at 0 °C for 45 min. Water (500 mL) and EtOAc (500 mL) were added and the phases were separated. The organic layer was successively washed with water (200 mL) and brine (200 mL), dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residual solid was recrystallised from MeOH to give the title compound (83.7 g, 80% yield) as yellow needles.

**b) 3-Amino-6-chloro-2-(ethylamino)pyridine**

A solution of SnCl<sub>2</sub>·H<sub>2</sub>O (616.3 g, 2.73 mol) in aqueous 12 N HCl (500 mL) was rapidly added to a solution of 6-chloro-2-(ethylamino)-3-nitropyridine (169.5 g, 0.84 mol) in AcOH (1.7 L) at room temperature. After 20 min, the mixture was cooled to 0 °C and water (250 mL) was added. Solid NaOH (240 g) was then added in small portions. The resulting suspension was filtered to remove tin salts. The filtrate was diluted with water (3.5 L), the solution rendered basic by addition of aqueous 10 N NaOH and then extracted with EtOAc (3 × 1.7 L). The combined organic layers were washed with brine (1 L), dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 3:2) to give the title compound (89.9 g, 62% yield) as a brown solid: MS (ESI) m/z 172/174 (MH)<sup>+</sup>.

**c) 5-Bromo-2-chloro-N-[2-(ethylamino)-6-chloro-3-pyridinyl]-3-pyridinecarboxamide**

A solution of 5-bromo-2-chloro-3-pyridinecarbonyl chloride (30.0 g, 97.0 mmol) in MeCN (100 mL) was added via cannula to a solution of 3-amino-6-chloro-2-(ethylamino)pyridine (16.6 g, 97.0 mmol) in MeCN (180 mL) containing solid NaHCO<sub>3</sub> (14.2 g, 169 mmol) at room temperature. The

mixture was stirred at room temperature for 1 h. Water (200 mL) was added and the mixture was stirred for 10 min. The resulting suspension was filtered. The solid was washed with Et<sub>2</sub>O (50 mL) and concentrated from pyridine solution (3 × 50 mL) to give the title compound (28.4 g, 75% yield).

5

**d) 8-Bromo-2-chloro-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

A 1 M solution of NaHMDS in THF (167.5 mL, 167.5 mmol) was slowly added to a solution of 5-bromo-2-chloro-N-[2-(ethylamino)-6-chloro-3-pyridinyl]-3-pyridinecarboxamide (28.4 g, 72.8 mmol) in pyridine (146 mL) heated to 50 °C. The reaction mixture was stirred at 50 °C for 1.5 h. The mixture was then poured into a mixture of water and ice (1 L) and, after 1 h, the resulting suspension was filtered. The solid was washed with water and dried under reduced pressure to give the title compound (23.4 g, 91% yield).

15

**e) 8-Bromo-2-chloro-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

Solid NaH (60% oil dispersion, 3.46 g, 86.1 mmol) was added over 30 min to a solution of 8-bromo-2-chloro-5,11-dihydro-11-ethyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (23.4 g, 66.3 mmol) in DMF (220 mL) at 50 °C. The mixture was stirred at 50 °C for 1 h then was allow to cool to room temperature. The mixture was poured into water (1 L) and the resulting suspension was filtered. The solid was successively washed with water and hexane then dried under reduced pressure to give the title compound (23.0 g, 94% yield).

25

**f) 2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

Allyltributyltin (21.3 mL, 68.7 mmol) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (3.61 g, 3.12 mmol) were added to a degassed (N<sub>2</sub> through solution for 30 min) solution of 8-bromo-2-chloro-5,11-dihydro-11-ethyl-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (23.0 g, 62.5 mmol) in DMF (312 mL). The mixture was

30

heated to 90 °C for 2 h. The mixture was concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 7:3) to give the title compound (13.4 g, 65% yield).

- 5 **g) 2-Chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**  
Ozonised oxygen was introduced into a cold (-78 °C) solution of 2-chloro-5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (13.4 g, 40.7 mmol) in MeOH (102 mL) and CH<sub>2</sub>Cl<sub>2</sub> (102  
10 mL) until complete disappearance of the alkene. Nitrogen was bubbled through the solution to remove excess O<sub>3</sub>. Solid NaBH<sub>4</sub> (2.69 g, 71.1 mmol) was next added in small portions and the mixture was allowed to warm slowly to room temperature. After 1 h, aqueous saturated NH<sub>4</sub>Cl (150 mL) was added and the mixture stirred for 20 min. The organic solvents were  
15 removed under reduced pressure. Water (100 mL) was added to the aqueous solution. The solution was extracted with CHCl<sub>3</sub> (3 × 200 mL). The combined organic layers were dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:CHCl<sub>3</sub>, 4:1) to give the title compound (10.4 g, 77% yield).

20

- h) 2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-[2-(4-quinolinylloxy)ethyl]-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**  
Diethyl azodicarboxylate (DEAD) (4.26 mL, 27.0 mmol) was added drop-  
wise to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-  
25 methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (6.00 g, 18.0 mmol), 4-hydroxyquinoline (3.92 g, 27.0 mmol) and Ph<sub>3</sub>P (7.09 g, 27.0 mmol) in THF (90 mL) at room temperature. The mixture was stirred at room temperature for 1h then was concentrated under reduced pressure. The residue was  
purified by flash chromatography (EtOAc:MeOH, 95:5) to give the title  
30 compound (6.22 g, 75% yield) as a white solid: MS (ESI) m/z 460/462.

**Example III (entry 6, Table 1)**

**2-Chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-(4-quinolinylloxy)-ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

5

**a) 6-Chloro-2-(cyclopropylamino)-3-nitropyridine**

A solution of cyclopropylamine (1.25 g, 22.0 mmol) in toluene (11 mL) was added over 10 min to an ice-cold solution of 2,6-dichloro-3-nitropyridine (2.00 g, 10.4 mmol) in toluene (10 mL). The mixture was stirred at 0 °C for 1 h and at room temperature for 2 h. Water (50 mL) was added to the mixture and the phases were separated. The organic layer was washed with brine (25 mL), dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 3:2) to give the title compound (1.97 g, 89% yield) as a yellow solid.

15

**b) 2-Chloro-11-cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

The title compound was prepared from 6-chloro-2-(cyclopropylamino)-3-nitropyridine using a method similar to that described for the 11-ethyl analog in example II.

20

**c) 2-Chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

Diethyl azodicarboxylate (DEAD) (85.6 µL, 0.54 mmol) was added dropwise to a solution of 2-chloro-11-cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.36 mmol), 4-hydroxyquinoline (78.9 mg, 0.54 mmol) and Ph<sub>3</sub>P (143 mg, 0.54 mmol) in THF (1.8 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5 µ, 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in

30



25 min) to give the trifluoroacetic acid salt of the title compound (17.7 g, 77% yield) as a white solid: MS (ESI) m/z 472/474 (MH)<sup>+</sup>.

**Example IV (entry 4, Table 1)**

5 ***5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one***

**a) 2,6-Difluoro-3-nitropyridine**

To a mixture of concentrated sulphuric acid (600 mL) and fuming nitric acid  
10 (90%, 400mL) in a ice-bath (internal temperature maintained at 5-10 °C) was added drop-wise 2,6-difluoropyridine (200 g, 1.74 mol). The resulting mixture was stirred overnight at room temperature. The mixture was poured slowly into 3 kg of ice and extracted with Et<sub>2</sub>O (2 × 2 L). The combined organic layers were washed with aqueous 1.5 N NaOH (2 × 1 L), then with  
15 aqueous saturated NaHCO<sub>3</sub> (400 mL) or until pH is around 8-9. The organic layers were dried over MgSO<sub>4</sub>, filtrated and concentrated under reduced pressure until constant weight (to remove unreacted 2,6-difluoropyridine: 10-12%). The title compound was obtained as a yellow liquid (207.3 g, 74% yield).

20

**b) 2-(Ethylamino)-6-fluoro-3-nitropyridine**

To a solution of 2,6-difluoro-3-nitropyridine (45.7 g, 285 mmol) in THF (500 mL) at -40 °C was added drop-wise a solution of ethylamine (25.7 g, 570 mmol) in THF (250 mL). After 30 min, the reaction mixture was concentrated  
25 under reduced pressure and the residue was dissolved in EtOAc. The organic phase was washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The resulting yellow solid was purified by flash chromatography (15% EtOAc in hexane) to give the title compound (43.2 g, 82% yield) as a yellow solid.

30

**c) 3-Amino-2-(ethylamino)-6-fluoropyridine**

A solution of 2-(ethylamino)-6-fluoro-3-nitropyridine (43.2 g, 230 mmol) in THF (1 L) was stirred overnight at room temperature under hydrogen (1 atm.) in the presence of 20% Pd(OH)<sub>2</sub>/C (4.35 g). The catalyst was removed  
5 by filtration through diatomaceous earth. The filtrate was concentrated under reduced pressure to give the title compound (36.3 g, 95% yield) as a black solid.

**d) 2-Chloro-N-[2-(ethylamino)-6-fluoro-3-pyridinyl]-5-bromo-3-pyridinecarboxamide**

To a cooled solution (4 °C) of 3-amino-2-(ethylamino)-6-fluoropyridine (31.0 g, 200 mmol) in MeCN (160 mL) was added solid NaHCO<sub>3</sub> (50.4 g, 600 mmol). After 15 min, a solution of 5-bromo-2-chloro-3-pyridinecarbonyl chloride (1 equiv., 200 mmol) in MeCN (155 mL) was added. After 60 min at  
15 room temperature, the reaction mixture was poured into water (1.2 L) and stirred for 30 min. The resulting solid was filtered, dried under reduced pressure at 50 °C overnight. The title compound (73.7 g, 99% yield) was obtained as a black solid.

**e) 8-Bromo-5,11-dihydro-11-ethyl-2-fluoro-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

To a solution of the 2-chloro-N-[2-(ethylamino)-6-fluoro-3-pyridinyl]-5-bromo-3-pyridinecarboxamide (73.5 g, 216 mmol) in pyridine (435 mL) at 50 °C was added drop-wise a 1 M solution of NaHMDS in THF (520 mL, 520  
25 mmol). After 10 min, the reaction was allowed to cool to room temperature, then poured over ice water (2 L). The resulting solid was filtered, rinse with water and then hexane. The solid was dried under reduced pressure to give the title compound (50.6 g, 69% yield) as a dark green solid.

**f) 8-Bromo-5,11-dihydro-11-ethyl-2-fluoro-5-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

To a solution of the 8-bromo-5,11-dihydro-11-ethyl-2-fluoro-6*H*-dipyrido[3,2-  
b:2',3'-e][1,4]diazepin-6-one (44 g, 130.5 mmol) in DMF (520 mL) was added  
NaH (4.28 g, 178 mmol), and the mixture was heated to 50 °C for 30 min. The  
reaction mixture was cooled to room temperature and treated with MeI (24.4  
5 mL, 522 mmol). After 1.5 h, the reaction mixture was poured over ice water.  
The solid was filtered, washed with water and then hexane, dried under  
reduced pressure to give the title compound (43.2 g, 94% yield) as dark grey  
solid.

10 **g) 5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-(2-propenyl)-6*H*-dipyrido[3,2-  
b:2',3'-e][1,4]diazepin-6-one**  
Allyltributyltin (32.0 mL, 103.4 mmol) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (5.43 g, 4.70 mmol)  
were added to a degassed (N<sub>2</sub> through solution for 45 min) solution of 8-  
bromo-5,11-dihydro-11-ethyl-2-fluoro-5-methyl-6*H*-dipyrido[3,2-b:2',3'-e]  
15 [1,4]diazepin-6-one (33.0 g, 94.0 mmol) in DMF (470 mL). Additional  
amounts of Pd(Ph<sub>3</sub>)<sub>4</sub> (1.09 g, 0.94 mmol added after 1, 2, 3, 4, 5 h of reaction)  
were added to complete the reaction. The mixture was heated to 90 °C for 6  
h. The mixture was concentrated under reduced pressure. The residue was  
purified by flash chromatography (hexanes:EtOAc, 8:2 to 7:3) to give the title  
20 compound (22.4 g, 76% yield).

**h) 5,11-Dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-5-methyl-6*H*-  
dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**  
A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of  
25 5,11-dihydro-11-ethyl-2-fluoro-5-methyl-8-(2-propenyl)-6*H*-dipyrido[3,2-  
b:2',3'-e][1,4]diazepin-6-one (22.38 g, 71.6 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and  
MeOH (100 mL) for 3 h. A stream of N<sub>2</sub> was next bubbled through the solution  
for 15 min and then solid NaBH<sub>4</sub> (5.05 g, 133 mmol) was added to the solution.  
The reaction mixture was allowed to warm to room temperature. After 1 h, an  
30 additional portion of NaBH<sub>4</sub> (1.62 g, 43.0 mmol) was added to the reaction

mixture. After an additional hour, aqueous saturated  $\text{NH}_4\text{Cl}$  (150 mL) was added and the mixture was stirred at room temperature for 30 min. The organic solvents were removed under reduced pressure. Water (200 mL) was added and the mixture was extracted with  $\text{CHCl}_3$  ( $3 \times 300$  mL). The combined  
5 organic layers were dried ( $\text{MgSO}_4$ ), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography ( $\text{EtOAc}:\text{CHCl}_3$ , 4:1) to give the title compound (19.7 g, 72% yield) as a white solid.

i) **5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**  
Diethyl azodicarboxylate (DEAD) (14.3 mL, 91.0 mmol) was added drop-  
wise to a solution of 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-5-  
methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (19.2 g, 60.7 mmol), 4-  
hydroxyquinoline (13.2 g, 91.0 mmol) and  $\text{Ph}_3\text{P}$  (23.9 g, 91.0 mmol) in THF  
15 (300 mL) at room temperature. The mixture was stirred at room temperature for 1h then was concentrated under reduced pressure. The residue was purified by flash chromatography ( $\text{EtOAc}:\text{MeOH}$ ; 95:5) to give the title compound (17.9 g, 67% yield) as a white solid: MS (ESI)  $m/z$  444 (MH)<sup>+</sup>.

20 **Example V (entry 1, Table 1)**  
**2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

a) **2-Chloro-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide**  
25  $\text{NaHCO}_3$  (21.72 g, 258.6 mmol) was added to a solution of 3-amino-2,6-dichloro-4-methylpyridine (41.61 g, 235.1 mmol; prepared as described by K. G. Grozinger *et al.* J. Heterocyclic Chem. 1995, 32, 259-263) in MeCN (350 mL) and the resulting suspension was stirred for 15 min. A solution of 2-chloronicotinyl chloride (43.44 g, 246.8 mmol) in MeCN (500 mL) was then  
30 introduced over 30 min. The resulting suspension was stirred at room temperature. After 24 h, the solution was found to be acidic and so an

additional amount of  $\text{NaHCO}_3$  (3.00 g, 35.7 mmol) was added. The suspension was then stirred at room temperature for an additional 2 days. The mixture was then poured into a mixture of water (2 L) and ice (200 g) and stirred for 20 min. The solid was filtered and washed with water (500 mL) and the resulting solid was then dried over  $\text{P}_2\text{O}_5$  under reduced pressure to give the title compound (62.55 g, 84% yield) as a white powder.

**b) *N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide**

A solution of 2-chloro-*N*-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (63.17 g, 194.0 mmol) and ethylamine (28.0 g, 583 mmol) in xylenes (250 mL) was stirred at 120 to 125 °C in a steel autoclave for 7 h. The resulting suspension was poured into water (1 L), stirred for 15 min and filtered. The residue was dissolved in EtOAc and washed with water (3 ×), brine and was dried over  $\text{MgSO}_4$ . The filtrate was extracted with EtOAc and the combined organic extracts were washed with water and brine and dried over  $\text{MgSO}_4$ . The two fractions were then combined and excess xylenes removed by co-distillation with benzene to give the title compound (61.98 g, pink solid) as the major component of a mixture of compounds. The material, not purified at this point, was used directly in the subsequent reaction.

**c) 5-Bromo-*N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide**

Solid KOAc (22.2 g, 229 mmol) was added to a solution of crude *N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide (61.98 g, < 190 mmol) in AcOH (600 mL). A solution of  $\text{Br}_2$  (9.7 mL, 191 mmol) was then added over 30 min. After stirring 30 min, additional amounts of KOAc and  $\text{Br}_2$  were introduced until TLC indicated that no starting material remained (a total of 11.2 g of KOAc and 5.91 mL of  $\text{Br}_2$ ). The solution was then poured into water (2 L) and stirred for 20 min. The solid was removed

by filtration and then dissolved in Et<sub>2</sub>O (1 L). The ethereal solution was washed with water (3 ×), aqueous saturated NaHCO<sub>3</sub>, brine then dried (MgSO<sub>4</sub>) to give a mixture containing, as the major component, the title compound (77.7 g, yellow foam). The material was used directly in the subsequent reaction.

**d) 8-Bromo-2-chloro-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A solution of crude 5-bromo-*N*-(2,6-dichloro-4-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide (77.7 g, mmol) in anhydrous pyridine (1.0 L) was heated to 50 °C. A 1 M solution of NaHMDS in THF (418 mL, 418 mmol) was then added dropwise and stirring was continued for an additional 15 min. After cooling to room temperature, water (50 mL) was added and the mixture concentrated to two-thirds volume on a rotary evaporator. The residue was then poured into water (3 L). Filtration gave the title compound (23.44 g) as a tan solid. The filtrate was extracted with EtOAc (3 ×) and the combined extracts were washed with water and brine then dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 9:1 or CHCl<sub>3</sub>) to give an additional amount of the title compound (14.06 g, for a total of 37.50 g, 50% yield for 3 steps).

**e) 2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A solution of 8-bromo-2-chloro-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (12.6 g, 34.3 mmol) in DMF (250 mL) was degassed under reduced pressure for 20 min. Pd(PPh<sub>3</sub>)<sub>4</sub> (775 mg, 0.7 mmol) was then added followed by allyltributyltin (12.5 mL, 41.1 mmol). After degassing under reduced pressure for 10 min, the mixture was heated to 100 °C for 7 h. The mixture was then concentrated under reduced

pressure. The residue was purified by flash chromatography (hexane:EtOAc, 19:1) to give the title compound (8.04 g, 71% yield) as a yellow solid.

**f) 2-Chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

5 A stream of O<sub>3</sub> was introduced into a cold (-78 °C) solution of 2-chloro-5,11-dihydro-11-ethyl-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (8.04 g, 24.4 mmol) and Sudan III in CH<sub>2</sub>Cl<sub>2</sub> (120 mL) and MeOH (120 mL). When the pink solution turned brown, O<sub>2</sub> was bubbled  
10 through the solution for 10 min. NaBH<sub>4</sub> (1.12 g, 29.4 mmol) was then added and the solution was allowed to warm to room temperature. After 30 min, an additional amount of NaBH<sub>4</sub> (500 mg) was added. After 1 h, aqueous saturated NH<sub>4</sub>Cl was then added and the mixture stirred for 20 min. The solution was concentrated under reduced pressure and extracted with  
15 CH<sub>2</sub>Cl<sub>2</sub> (3 ×). The combined organic extracts were washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CHCl<sub>3</sub>:EtOH, 97:3) to give the title compound (6.01 g, 74% yield) as a white solid.

**g) 2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-[2-(4-quinolinyloxy)ethyl]-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

20 Diethyl azodicarboxylate (DEAD) (83 µL, 0.53 mmol) was added drop-wise to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.38 mmol), 4-  
25 hydroxyquinoline (76.3 mg, 0.53 mmol) and Ph<sub>3</sub>P (138 mg, 0.53 mmol) in THF (2.5 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 1 N aqueous NaOH (3 × 10 mL) and brine (15 mL) then was  
30 dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The

residue was purified by flash chromatography (EtOAc:MeOH, 9:1) to give the title compound (52 mg, 30% yield) as a white solid: MS (ESI) m/z 460/462 (MH)<sup>+</sup>.

5 **Example VI (entry 7, Table 1)**

**5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

- 10 **a) 8-Bromo-5,11-dihydro-11-ethyl-2-[(4-methoxyphenyl)methyl]amino}-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**  
8-Bromo-2-chloro-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (2.50 g, 6.80 mmol) was dissolved in dioxane (10 mL) in a glass pressure bottle and 4-methoxybenzylamine (4.0 mL, 30.6 mmol) was added to the solution. The mixture was heated to 170 °C for 5 days. The  
15 reaction mixture was cooled to room temperature and concentrated under reduced pressure. The residue was taken up in EtOAc and washed with 10% aqueous NH<sub>4</sub>Cl, dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure to give a brown oil which was chromatographed over silica gel (hexane:EtOAc, 4:1). The title compound was obtained as a yellow solid  
20 (1.35 g, 42% yield).

- b) 2-Amino-8-bromo-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**  
8-Bromo-5,11-dihydro-11-ethyl-2-[(4-methoxyphenyl)methyl]amino}-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (6.52 g, 18.7 mmol) was  
25 dissolved in trifluoroacetic acid (100 mL) and the solution was stirred magnetically at room temperature for 18 h. Trifluoroacetic acid was evaporated under reduced pressure and the residue was poured into a mixture containing water (450 mL), concentrated NH<sub>4</sub>OH (50 mL) and  
30 EtOAc (400 mL). The mixture was stirred magnetically for 2 h at room temperature. The organic phase was concentrated under reduced pressure.



The yellow precipitate was filtered off and washed with water, dried under reduced pressure to give 7.51 g of a yellow solid which was triturated in Et<sub>2</sub>O to give the title compound (5.12 g, 105% yield) as a yellow solid.

5    **c) 8-Bromo-5,11-dihydro-11-ethyl-2-fluoro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A plastic bottle was charged with 2-amino-8-bromo-5,11-dihydro-11-ethyl-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (5.10 g, 14.6 mmol). HF·pyridine (100 g) was added and the resulting suspension was cooled to 0°  
10    C. Sodium nitrite (1.12 g, 16.1 mmol) was added in several portions over 30 min to produce a purple solution. The mixture was then stirred for 16 h at room temperature. The reaction mixture was poured onto ice and 6 N NaOH. The beige suspension was extracted with EtOAc. The organic extracts were combined, dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was  
15    triturated with Et<sub>2</sub>O/hexane to give the title compound (4.30 g, 84% yield).

**d) 5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

Allyltributyltin (7.77 mL, 25.1 mmol) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (1.32 g, 1.14 mmol) were  
20    added to a degassed (N<sub>2</sub> through solution for 30 min) solution of 8-bromo-5,11-dihydro-11-ethyl-2-fluoro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (8.0 g, 22.8 mmol) in DMF (114 mL). The mixture was heated to 90 °C for 3 h. The mixture was poured into water (250 mL) and extracted with EtOAc (4 × 250 mL). The combined organic layers were  
25    washed with 20% aqueous NH<sub>4</sub>OH (250 mL) and brine (250 mL), dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexanes:EtOAc, 4:1 to 1:1) to give the title compound (3.16 g, 51% yield).

30    **e) 5,11-Dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

- A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of 5,11-dihydro-11-ethyl-2-fluoro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (3.16 g, 10.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (25 mL) and MeOH (25 mL) for 45 min. A stream of N<sub>2</sub> was next bubbled through the solution for 15 min and then solid NaBH<sub>4</sub> (669 mg, 17.1 mmol) was added to the solution. The reaction mixture was allowed to warm to room temperature. After 2 h, aqueous saturated NH<sub>4</sub>Cl (50 mL) was added and the mixture was stirred at room temperature for 15 min. The organic solvents were removed under reduced pressure. The residual aqueous solution was extracted with CHCl<sub>3</sub> (3 × 100 mL). The combined organic layers were dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:CHCl<sub>3</sub>, 1:1) to give the title compound (2.40 g, 75 % yield) as a white solid.
- f) **5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-(4-quinolinyloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**  
Diisopropyl azodicarboxylate (DIAD) (117 µL, 0.59 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (125 mg, 0.40 mmol), 4-hydroxyquinoline (86.0 mg, 0.59 mmol) and Ph<sub>3</sub>P (156 mg, 0.59 mmol) in THF (1.9 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (40 mL) and the solution was successively washed with 3 N aqueous NaOH (3 × 20 mL), aqueous saturated NH<sub>4</sub>Cl (15 mL), brine (15 mL) then was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5 µ, 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min) to give the

trifluoroacetic acid salt of the title compound (106 mg, 48% yield) as a white solid: MS (ESI)  $m/z$  444 (MH)<sup>+</sup>.

**Example VII (entry 5, Table 1)**

**5 11-(Cyclopropyl)-5,11-dihydro-2-fluoro-4-methyl-8-(2-(4-quinolinyl-*oxy*)-ethyl)-6H-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one**

**a) 11-Cyclopropyl-5,11-dihydro-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one**

10 The title compound was prepared from 8-bromo-2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one using a procedure similar to that of the corresponding 11-ethyl analog described in example VI.

15 **b) 11-Cyclopropyl-5,11-dihydro-2-fluoro-4-methyl-8-(2-(4-quinolinyl-*oxy*)ethyl)-6H-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one**  
Diisopropyl azodicarboxylate (DIAD) (131  $\mu$ L, 0.66 mmol) was added dropwise to a solution of 11-(cyclopropyl)-5,11-dihydro-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one (140  
20 mg, 0.44 mmol), 4-hydroxyquinoline (96.4 mg, 0.66 mmol) and Ph<sub>3</sub>P (124 mg, 0.66 mmol) in THF (2.2 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 3 N aqueous NaOH (3  $\times$  20 mL), aqueous  
25 saturated NH<sub>4</sub>Cl (20 mL) and brine (20 mL) then was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC (CombiPrep ADS-AQ 50  $\times$  20 mm, 5  $\mu$ , 120 A, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min) to give the  
30 trifluoroacetic acid salt of the title compound (104 mg, 43% yield) as a white solid: MS (ESI)  $m/z$  456 (MH)<sup>+</sup>.

**Example VIII (entry 8, Table 1)**

***5,11-Dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

5

**a) 5,11-Dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A mixture of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (168 mg, 0.51 mmol),  
10 ammonium formate (318 mg, 5.1 mmol) and 10% Pd/C (50 mg) in EtOH (6 mL) was stirred at room temperature. After 24 h, additional amounts of ammonium formate (200 mg, mmol) and 10% Pd/C (50 mg) were added. After stirring an additional 24 h, the reaction was concentrated under reduced pressure. The residue was dissolved in CHCl<sub>3</sub> and the mixture was  
15 washed with aqueous saturated NaHCO<sub>3</sub>, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure to give the title compound (122 mg, 81%) as a yellow oil.

**b) 5,11-Dihydro-11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**  
20 Diisopropyl azodicarboxylate (DIAD) (42 µL, 0.21 mmol) was added dropwise to a solution of 5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (42.0 mg, 0.14 mmol), 4-hydroxyquinoline (31 mg, 0.21 mmol) and Ph<sub>3</sub>P (55.4 mg, 0.21 mmol) in THF  
25 (0.7 mL) at room temperature. The mixture was stirred at room temperature for 2 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 3 N aqueous NaOH (3 × 20 mL), aqueous saturated NH<sub>4</sub>Cl (20 mL) and brine (20 mL) then was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced  
30 pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The solid was further purified by reverse phase HPLC

(CombiPrep ADS-AQ 50 × 20 mm, 5  $\mu$ , 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min) to give the trifluoroacetic acid salt of the title compound (14.2 mg, 19% yield) as a white solid: MS (ESI)  $m/z$  426 (MH)<sup>+</sup>.

5 **Example IX (entry 3, Table 1)**

**11-(Cyclopropyl)-5,11-dihydro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

10 **a) 5-Bromo-2-chloro-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide**

3-Amino-2,6-dichloro-4-methylpyridine (0.51 g, 2.85 mmol) was dissolved in toluene (35 mL) and pyridine (0.27 mL, 3.28 mmol) was added. 5-Bromo-2-chloro-3-pyridinecarbonyl chloride (0.80 g, 3.14 mmol) was then added dropwise over 30 min. The resulting mixture was stirred at room  
15 temperature for 1 h, diluted with water and extracted with toluene (2 ×). The combined organic extracts were dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The resulting thick oil was triturated with CH<sub>2</sub>Cl<sub>2</sub>, and the white solid collected via suction filtration to give the title compound (0.41 g, 36% yield).

20

**b) 5-Bromo-2-(cyclopropylamino)-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide**

A solution of 5-bromo-2-chloro-N-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (3.00 g, 7.61 mmol) and cyclopropylamine (2.11 mL,  
25 30.5 mmol) in xylenes was heated in a sealed tube at 120 °C for 24 h. The reaction was cooled to 0 °C, diluted with water and extracted with EtOAc (2 ×). The combined organic extracts were dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The resulting residue was purified by trituration (Et<sub>2</sub>O:hexane, 3:1) to give the title compound (2.35 g, 75% yield) as  
30 a pale yellow solid.

**c) 8-Bromo-2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A solution of 5-bromo-2-(cyclopropylamino)-*N*-(2,6-dichloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (105 mg, 0.254 mmol) in pyridine (3.0 mL) was heated to 50 °C while a 1 M solution of NaHMDS in THF (0.53 mL, 0.53 mmol) was added. The resulting solution was stirred at 50 °C for 3 h. The reaction was quenched with aqueous saturated NH<sub>4</sub>Cl, diluted with water and extracted with EtOAc (3 ×). The combined organic extracts were washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The resulting residue was purified via flash chromatography (hexane:EtOAc, 7:3) to give the title compound (32 mg, 33% yield).

**d) 2-Chloro-11-cyclopropyl-5,11-dihydro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A solution of 8-bromo-2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (0.046 g, 0.122 mmol) in DMF (2.0 mL) was degassed by sparging with N<sub>2</sub> for 10 min. Pd(PPh<sub>3</sub>)<sub>4</sub> (2.8 mg, 0.003 mmol) and allyltributyltin (0.45 mL, 0.146 mmol) were added and the resulting solution was heated to 90 °C for 2.5 h. The reaction mixture was diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 ×) and EtOAc (2 ×). The combined organic extracts were dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The resulting residue was purified via flash chromatography (hexane to hexane:EtOAc, 7:3) to give the title compound (0.033 g, 80% yield).

**e) 2-Chloro-11-cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A cold (-78 °C) solution of 2-chloro-11-cyclopropyl-5,11-dihydro-4-methyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (0.82 g, 2.43 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7.5 mL) and MeOH (7.5 mL) was saturated with O<sub>3</sub> until the solution turned pale blue. The reaction mixture was left at -78 °C for 10

min, then NaBH<sub>4</sub> (0.23 g, 9.71 mmol) was added and the mixture was allowed to warm slowly to room temperature over 2 h. The reaction was quenched with aqueous 10% citric acid solution and extracted with EtOAc (3 ×). The organic extracts were combined, washed with brine, dried (MgSO<sub>4</sub>),  
5 filtered and concentrated under reduced pressure. The resulting residue was purified via flash chromatography (MeOH:CH<sub>2</sub>Cl<sub>2</sub>, 2 to 10%) to give the title compound (0.41 g, 60% yield) as a white solid.

**f) 11-Cyclopropyl-5,11-dihydro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

10 The title compound was prepared using a method similar to that described for the 11-ethyl analog in example VIIIa.

**g) 11-Cyclopropyl-5,11-dihydro-4-methyl-8-[2-(4-quinolinylloxy)ethyl]-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

15 Diisopropyl azodicarboxylate (DIAD) (167 µL, 0.85 mmol) was added dropwise to a solution of 11-(cyclopropyl)-5,11-dihydro-8-(2-hydroxyethyl)-4-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (175 mg, 0.57 mmol), 4-hydroxyquinoline (123 mg, 0.85 mmol) and Ph<sub>3</sub>P (223 mg, 0.85 mmol) in THF  
20 (2.8 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (60 mL) and the solution was successively washed with 3 N aqueous NaOH (3 × 20 mL), aqueous saturated NH<sub>4</sub>Cl (20 mL) and brine (20 mL) then was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced  
25 pressure. The residue was purified by flash chromatography (EtOAc:MeOH; 95:5) to give the title compound (74 mg, 30% yield) as a white solid: MS (ESI) m/z 438 (MH)<sup>+</sup>.

**Example X (entry 21, Table 1)**

***5,11-Dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

- 5 Solid mCPBA (80-85%, *m*-chloroperbenzoic acid) (2.21 g, 10.2 mmol) was added to a solution of 5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (2.56 g, 6.02 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (30 mL) and THF (30 mL) at room temperature (note that the reaction can be performed in CH<sub>2</sub>Cl<sub>2</sub> alone). The mixture was stirred  
10 at room temperature for 1 h then was concentrated under reduced pressure. The residue was dissolved in CHCl<sub>3</sub> (400 mL), the solution was successively washed with aqueous 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (3 × 75 mL), aqueous saturated NaHCO<sub>3</sub> (3 × 75 mL) and brine (75 mL), dried (MgSO<sub>4</sub>), filtered and concentrated  
under reduced pressure. The residue was purified by flash chromatography  
15 (CHCl<sub>3</sub>:MeOH, 9:1) to give the title compound (2.01 g, 76% yield) as a white solid: MS (ESI) *m/z* 442 (MH)<sup>+</sup>.

**Example XI (entry 17, Table 1)**

20 ***2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

- Using a procedure identical to that of example X, 2-chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e]  
[1,4]diazepin-6-one (2.70 g, 5.87 mmol) gave the title compound (2.05 g, 73%  
25 yield) as a white solid: MS (ESI) *m/z* 476/478 (MH)<sup>+</sup>.

**Example XII (entry 11, Table 1)**

30 ***5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

- Solid mCPBA (457 mg, 2.12 mmol) was added to a solution of 5,11-dihydro-2-fluoro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (470 mg, 1.06 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The



mixture was stirred at room temperature for 13 h then was concentrated under reduced pressure. The residue was dissolved in  $\text{CHCl}_3$  (75 mL), the solution was successively washed with aqueous 10%  $\text{Na}_2\text{S}_2\text{O}_3$  ( $3 \times 25$  mL), aqueous saturated  $\text{NaHCO}_3$  ( $3 \times 25$  mL) and brine (25 mL), dried ( $\text{MgSO}_4$ ),  
5 filtered and concentrated under reduced pressure. The residue was purified by flash chromatography ( $\text{CHCl}_3$ :MeOH, 19:1 to 9:1) to give the title compound (250 mg, 51% yield) as a white solid: MS (ESI)  $m/z$  460 ( $\text{MH}^+$ ).

**Example XIII (entry 10, Table 1)**

10 ***2-Chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-[(1-oxido-4-quinolin-yl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

Solid mCPBA (177 mg, 0.82 mmol) was added to a solution of 2-chloro-11-cyclopropyl-5,11-dihydro-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-  
15 dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (143 mg, 0.30 mmol) in  $\text{CH}_2\text{Cl}_2$  (3 mL) and THF (3 mL). The mixture was stirred at room temperature for 14 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (50 mL), the solution was successively washed with aqueous 10%  $\text{Na}_2\text{S}_2\text{O}_3$  ( $3 \times 10$  mL), aqueous saturated  $\text{NaHCO}_3$  ( $3 \times 10$  mL) and brine (10  
20 mL), dried ( $\text{MgSO}_4$ ), filtered and concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH, 95:5 to  $\text{CH}_2\text{Cl}_2$ :MeOH, 9:1). The impure compound was further purified by reverse phase HPLC (CombiPrep ADS-AQ  $50 \times 20$  mm,  $5 \mu$ , 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min). The pure  
25 fractions were treated with aqueous saturated  $\text{NaHCO}_3$ , extracted with EtOAc and the solution was dried ( $\text{MgSO}_4$ ), filtered and concentrated under reduced pressure to give the title compound (28.3 mg, 19%) as a white solid: MS (ESI)  $m/z$  488/490 ( $\text{MH}^+$ ).

**Example XIV (entry 14, Table 1)**

***2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-[(1-oxido-4-quinolinyl)-oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

- 5 Solid mCPBA (46.9 mg, 0.22 mmol) was added to a solution of 2-chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (50.0 mg, 0.11 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.2 mL). The mixture was stirred at room temperature for 1 h then was concentrated under reduced pressure. The residue was dissolved in EtOAc (50 mL), the  
10 solution was successively washed with aqueous 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (3 × 10 mL), aqueous saturated NaHCO<sub>3</sub> (3 × 10 mL) and brine (10 mL), dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by reverse phase HPLC (CombiPrep ADS-AQ 50 × 20 mm, 5 μ, 120 Å, 5-100% MeCN + 0.10% TFA / water + 0.10% TFA in 25 min). The pure  
15 fractions were treated with aqueous saturated NaHCO<sub>3</sub>, extracted with EtOAc and the solution was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure to give the title compound (32.5 mg, 63%) as a white solid: MS (ESI) m/z 488/490 (MH)<sup>+</sup>.

20 **Example XV (entry 9, Table 1)**

***5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-[(1-oxido-4-quinolinyl)-oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

- Using a procedure similar to that of example XIV, 5,11-dihydro-11-ethyl-2-  
25 fluoro-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.28 mmol) gave the title compound (40 mg, 31%) as a white solid: MS (ESI) m/z 460 (MH)<sup>+</sup>.

**Example XVI (entry 12, Table 1)**

- 30 ***5,11-Dihydro-11-ethyl-4-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

Solid mCPBA (163 mg, 0.75 mmol) was added to a solution of 5,11-dihydro-

11-ethyl-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e]  
[1,4]diazepin-6-one (144 mg, 0.34 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7.5 mL). The solution  
was stirred at room temperature for 2 h. The mixture was diluted with  
CH<sub>2</sub>Cl<sub>2</sub> (125 mL), washed successively with aqueous 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (2 × 25  
5 mL), aqueous saturated NaHCO<sub>3</sub> (2 × 25 mL) and brine (25 mL), then was  
dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The  
residue was purified by flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:MeOH, 10:1) to give  
the title compound (102 mg, 68% yield) as a white solid: MS (ESI) m/z 442  
(MH).

10

**Example XVII (entry 15, Table 1)**

**11-Cyclopropyl-5,11-dihydro-11-ethyl-4-methyl-8-{2-[(1-oxido-4-quinolin-  
yl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

15 Using a procedure similar to that of example XIV, 11-cyclopropyl-5,11-  
dihydro-11-4-methyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e]  
[1,4]diazepin-6-one (37.1 mg, 0.085 mmol) gave the title compound (35 mg,  
91%) as a white solid: MS (ESI) m/z 454 (MH)<sup>+</sup>.

20 **Example XVIII (entry 25, Table 1)**

**2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-(5-quinolinylloxy)ethyl}-6H-  
dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

Diethyl azodicarboxylate (DEAD) (77 µL, 0.49 mmol) was added drop wise  
25 to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-  
6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (125 mg, 0.38 mmol), 5-  
hydroxyquinoline (70.9 mg, 0.49 mmol) and Ph<sub>3</sub>P (128 mg, 0.49 mmol) in  
THF (2.5 mL) at room temperature. After 1.5 h, additional amounts of DEAD  
(35 µL, 0.22 mmol) and PPh<sub>3</sub> (59 mg, 0.22 mmol) were added to the mixture.  
30 The mixture was stirred at room temperature for 1 h then was diluted in  
EtOAc (50 mL) and the solution was successively washed with aqueous 1 N

NaOH (4 × 10 mL) and brine (15 mL) then was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:acetone, 85:15). The resulting solid was triturated with Et<sub>2</sub>O to give the title compound (38 mg, 22% yield) as a white solid: MS (ESI) m/z 460/462 (MH)<sup>+</sup>.

**Example XIX (entry 20, Table 1)**

***2-Chloro-5,11-dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

Diethyl azodicarboxylate (DEAD) (160 µL, 1.01 mmol) was added drop-wise to a solution of 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (225 mg, 0.68 mmol), 5-hydroxyquinoline (147 mg, 1.01 mmol) and Ph<sub>3</sub>P (266 mg, 1.01 mmol) in THF (3.4 mL) at room temperature. The mixture was stirred at room temperature for 3 h then was concentrated under reduced pressure. The residue was partially purified by flash chromatography (EtOAc:MeOH; 95:5). The fractions containing the 5-quinolinyl-oxy derivative were concentrated, dissolved in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) and THF (5 mL) and treated with mCPBA (80%, 248 mg, 1.15 mmol) at room temperature. The solution was stirred at room temperature for 2.5 h, then was concentrated under reduced pressure. The residue was dissolved in EtOAc (50 mL), washed successively with aqueous 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (3 × 10 mL), aqueous saturated NaHCO<sub>3</sub> (3 × 10 mL) and brine (10 mL), then was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH, 95:5 to CH<sub>2</sub>Cl<sub>2</sub>:MeOH, 9:1) to give the title compound (113 mg, 35% yield for 2 steps) as a white solid: MS (ESI) m/z 476/478 (MH)<sup>+</sup>.

**Example XX (entry 18, Table 1)**

***5,11-Dihydro-11-ethyl-2-fluoro-5-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one***

Using a procedure identical to that of example XIX, 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-5-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (267 mg, 0.84 mmol) gave the title compound (204 mg, 52% yield for 2 steps) as a white solid: MS (ESI) *m/z* 460 (MH)<sup>+</sup>.

**Example XXI (entry 19, Table 1)**

***2-Chloro-5,11-dihydro-11-ethyl-4-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one***

Using a procedure similar to that of example XIX but using diisopropyl azodicarboxylate (DIAD) instead of diethyl azodicarboxylate, 2-chloro-5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (223 mg, 0.67 mmol) gave the title compound (43.4 mg, 14% yield for 2 steps) as a white solid: MS (ESI) *m/z* 476/478 (MH)<sup>+</sup>.

**Example XXII (entry 23, Table 1)**

***5,11-Dihydro-11-ethyl-2-fluoro-4-methyl-8-{2-[(1-oxido-5-quinolinyl)-oxy]ethyl}-6H-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one*** Using a procedure

similar to that of example XXI, 5,11-dihydro-11-ethyl-2-fluoro-8-(2-hydroxyethyl)-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (250 mg, 0.79 mmol) gave the title compound (64.7 mg, 18% yield for 2 steps) as a white solid: MS (ESI) *m/z* 460 (MH)<sup>+</sup>.

**Example XXIII, Entry 22, Table 1**

***2-Chloro-5,11-dihydro-11-ethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one***

MS (ESI) *m/z* 462/464 (MH)<sup>+</sup>.

**Example XXIV, Entry 26, Table 1**

***5,11-Dihydro-5,11-diethyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one***

MS (ESI) *m/z* 440 (MH)<sup>+</sup>.

**Example XXV, Entry 27, Table 1**

**5,11-Dihydro-5,11-diethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one** MS (ESI) m/z 456 (MH)<sup>+</sup>.

5

**Example XXVI (Entry 28, Table 1)**

**2,5-Dimethyl-5,11-dihydro-11-ethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

- 10 **a) 5-Bromo-2-chloro-N-(6-chloro-2-methyl-3-pyridinyl)-3-pyridinecarboxamide**  
NaHCO<sub>3</sub> (3.9 g, 46.4 mmol) was added to a solution of 3-amino-2-chloro-6-methylpyridine (2.2 g, 15.4 mmol; prepared as described by K. G. Grozinger et al. J. Heterocyclic Chem. 1995, 32, 259-263) in MeCN (50 mL). The  
15 resulting suspension was stirred for 15 min. and a solution of crude 5-bromo-2-chloro-3-pyridinecarbonyl chloride (prepared from 5-bromo-2-hydroxy-3-pyridinecarboxylic acid and SOCl<sub>2</sub> [as described by T. W. Gero et al. in Synth. Commun. 1989, 19, 553-559 (incorporated herein by reference) but with omission of the aqueous work-up] (4 g) in MeCN (10 mL) was  
20 introduced over 30 min. The resulting suspension was stirred at room temperature. After 24 h, the mixture was poured into a mixture of water (100 mL) and ice (10 g) and stirred for 20 min. The suspension was filtered and the resulting solid was washed with water (50 mL) and hexane (25 mL), then dried over P<sub>2</sub>O<sub>5</sub> under reduced pressure to give the title compound (1.8 g,  
25 31% yield) as a white powder.

**b) 5-Bromo-N-(2-chloro-6-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide**

- A solution of 5-bromo-2-chloro-N-(2-chloro-6-methyl-3-pyridinyl)-3-pyridinecarboxamide (1.7 g, 4.8 mmol) and ethylamine (2M in THF, 10.5 mmol, 5.2 mL) in THF (5 mL) was stirred at 90 to 95 °C in a steel autoclave

for 16 h. The resulting mixture was poured into water (100 mL), stirred for 15 min and filtered. The solid was washed with water followed by hexane to give the title compound as a yellow solid (1.58 g, 89% yield).

5    **c) 8-Bromo-5,11-dihydro-2,5-dimethyl-11-ethyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

A solution of crude 5-bromo-*N*-(2-dichloro-6-methyl-3-pyridinyl)-2-(ethylamino)-3-pyridinecarboxamide (0.7 g, 1.8 mmol) in anhydrous pyridine (18 mL) was heated to 50 °C. A 1 M solution of NaHMDS in THF (7.2 mL, 7.2 mmol) was then added dropwise and stirring was continued for an additional 3h. After cooling to room temperature, methyl iodide (0.6 mL, 9 mmol) was added and the reaction was stirred overnight. Water (25 mL) was added and the mixture was extracted with EtOAc (3 ×) and the combined extracts were washed with water and brine then dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 8:2) to give the title compound (259 mg, 41% yield).

20    **d) 5,11-Dihydro-2,5-dimethyl-11-ethyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

A solution of 8-bromo-5,11-dihydro-2,5-dimethyl-11-ethyl-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (258 mg, 0.7 mmol) in DMF (7.4 mL) was degassed under reduced pressure for 20 min. Pd(PPh<sub>3</sub>)<sub>4</sub> (43 mg, 0.04 mmol) was then added followed by allyltributyltin (0.3 mL, 0.85 mmol). After 25 degassing under reduced pressure for 10 min, the mixture was heated to 100 °C for 1.5 h. The mixture was then concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 8:2) to give the title compound (184 mg, 98% yield) as a yellow solid.

**e) 5,11-Dihydro-2,5-dimethyl-11-ethyl-8-(2-hydroxyethyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

A stream of O<sub>3</sub> was introduced into a cold (-78 °C) solution of 5,11-dihydro-2,5-dimethyl-11-ethyl-8-(2-propenyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (187 mg, 0.6 mmol) and Sudan III in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) and MeOH (3 mL). When the pink solution turned brown, O<sub>2</sub> was bubbled through the solution for 10 min. NaBH<sub>4</sub> (57 mg, 1.52 mmol) was then added and the solution was allowed to warm to room temperature. After 30 min, an additional amount of NaBH<sub>4</sub> (20 mg) was added. After 2 h, aqueous saturated NH<sub>4</sub>Cl was then added and the mixture stirred for 20 min. The solution was concentrated under reduced pressure and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x). The combined organic extracts were washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:Hexane 6:4) to give the title compound (111 mg, 58% yield) as a white solid.

**f) 2,5-Dimethyl-5,11-dihydro-11-ethyl-8-[2-(4-quinolinylloxy)ethyl]-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

Diethyl azodicarboxylate (DEAD) (139 µL, 0.88 mmol) was added dropwise to a solution of 5,11-dihydro-2,5-dimethyl-11-ethyl-8-(2-hydroxyethyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (110 mg, 0.35 mmol), 4-hydroxyquinoline (128 mg, 0.88 mmol) and Ph<sub>3</sub>P (231 mg, 0.88 mmol) in THF (3.5 mL) at room temperature. The mixture was stirred at room temperature for 16h. The reaction mixture was diluted in EtOAc (60 mL) and the solution was successively washed with 1 N aqueous NaOH (3 × 10 mL) and brine (15 mL) then was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc then EtOAc:MeOH, 9.5:0.5) to give the title compound (57 mg, 37% yield) as a white solid: MS (ESI) m/z 440 (MH)<sup>+</sup>.



g) **5,11-Dihydro-2,5-dimethyl-11-ethyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**  
Solid mCPBA (80-85%, m-chloroperbenzoic acid) (60 mg, 0.28 mmol) was

- 5 added to a solution of 5,11-dihydro-2,5-dimethyl-11-ethyl-8-{2-(4-quinolinylloxy)ethyl}-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one (55 mg, 0.13 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1.3 mL) at room temperature. The mixture was stirred at room temperature for 2.5 h then diluted with CH<sub>2</sub>Cl<sub>2</sub>. The resulting solution was successively washed with aqueous 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, aqueous  
10 saturated NaHCO<sub>3</sub> and brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (MeOH/EtOAc, 2% to 5%) to give the title compound (59 mg, 99% yield) as a white solid: MS (ESI) m/z 456 (MH)<sup>+</sup>.

- 15 **Example XXVII (Entry 29, Table 1)**  
**5,11-Dihydro-11-ethyl-5-methyl-8-{2-[(1-oxido-4-quinolinyl)oxy]ethyl}-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e] [1,4]diazepin-6-one**

a) **2-(Ethylamino)-3-nitro-6-(trifluoromethyl)pyridine**

- 20 To a solution of 2-nitroacetamide ammonium salt (4 g, 33 mmol) in water (165 mL) was added piperidinium acetate (33 mmol) in water followed by slow addition of 4-ethoxy-1,1,1-trifluoro-3-buten-2-one (6.1 mL, 43 mmol) in MeOH (11 mL). The reaction was stirred for 2 h at room temperature and at reflux for 3 h. The reaction mixture was allowed to cool to 45 °C and aqueous  
25 HCl (1N) was added until the pH was acidic. After 1 h at room temperature, the reaction mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x). The combined organic layer was successively washed with water and brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc, then EtOAc:MeOH, 9:1) to give 3-nitro-6-  
30 (trifluoromethyl)-2(1H)-pyridinone (2.7 g, 40% yield) as a yellow solid.

To a solution of 3-nitro-6-(trifluoromethyl)-2(1*H*)-pyridinone (2.4 g, 11.5 mmol) in DMF (69 mL) was added sodium hydride (350 mg, 13.8 mmol). The reaction was stirred for 30 min at 45 °C, cooled to room temperature and *N*-phenyltrifluoromethanesulfonimide (4.9 g, 13.8 mmol) was added. After 1 h, a  
5 2 M solution of ethylamine in THF (13 mL, 25 mmol) was added. The reaction was stirred at room temperature overnight. The reaction mixture was poured into water (100 mL) and extracted with EtOAc (3x). The combined organic layer was successively washed water and brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was  
10 purified by flash chromatography (hexane:EtOAc, 9:1) to give the desired compound (1.1 g, 41% yield) as a yellow oil.

**b) 3-Amino-2-(ethylamino)-6-(trifluoromethyl)pyridine**

A solution of 2-(ethylamino)-3-nitro-6-(trifluoromethyl)pyridine (1.1 g, 4.9  
15 mmol) in MeOH (40 mL) was stirred overnight at room temperature under hydrogen (1 atm.) in the presence of 20% Pd(OH)<sub>2</sub>/C (100 mg). The catalyst was removed by filtration through diatomaceous earth. The filtrate was concentrated under reduced pressure to give the title compound as an orange oil (1 g).

20

**c) 2-Chloro-*N*-[2-(ethylamino)-6-(trifluoromethyl)-3-pyridinyl]-5-bromo-3-pyridinecarboxamide**

To a cooled solution of 3-amino-2-(ethylamino)-6-trifluoromethyl pyridine (1 g, 4.9 mmol) in MeCN (24 mL) was added solid NaHCO<sub>3</sub> (906 mg, 11 mmol).  
25 After 5 min, crude 5-bromo-2-chloro-3-pyridinecarbonyl chloride (prepared from 5-bromo-2-hydroxy-3-pyridinecarboxylic acid and SOCl<sub>2</sub> [as described by T. W. Gero et al. in Synth. Commun. 1989, 19, 553-559 (incorporated herein by reference) but with omission of the aqueous work-up] was added (1 equiv., 4.9 mmol). After 2 h, the reaction mixture was poured over

ice/H<sub>2</sub>O (1.5 L) and the resulting solid was filtered, rinsed with H<sub>2</sub>O and then hexane. After drying under reduced pressure overnight, the title compound was obtained as a light brown solid (1.6 g, 77% yield).

5    **d) 8-Bromo-5,11-dihydro-11-ethyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

To a solution of 2-chloro-*N*-[2-(ethylamino)-6-(trifluoromethyl)-3-pyridinyl]-5-bromo-3-pyridinecarboxamide (766 mg, 1.8 mmol) in DMF (18 mL) was added NaH (137 mg, 5.4 mmol). The reaction was stirred at 80 °C. After 1 h  
10    the reaction was allowed to cool to room temperature, and then was poured in water (20 mL). The resulting mixture was extracted with EtOAc (3x). The combined organic layer was successively washed with water and brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 7:3) to give the desired  
15    compound (200 mg, 28% yield) as a yellow solid.

**e) 8-Bromo-5,11-dihydro-11-ethyl-5-methyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

To a solution of the 8-bromo-5,11-dihydro-11-ethyl-2-(trifluoromethyl)-6H-  
20    dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (200 mg, 0.52 mmol) in DMF (2.6 mL) was added NaH (20 mg, 0.78 mmol), and the mixture was heated to 50 °C for 30 min. The reaction mixture was cooled to room temperature and treated with MeI (97 µL, 1.56 mmol). After 16 h, the reaction mixture was diluted with water. The resulting mixture was extracted with EtOAc (3x).  
25    The combined organic layer was successively washed with water and brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 8.5:1.5) to give the title compound (110 mg 53% yield) as a white foam.

**f) 5,11-Dihydro-11-ethyl-5-methyl-8-(2-propenyl)-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

Allyltributyltin (98  $\mu$ L, 0.32 mmol) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (16 mg, 0.02 mmol) were added to a degassed (N<sub>2</sub> through solution for 30 min) solution of 8-bromo-5,11-dihydro-11-ethyl-5-methyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (11 mg, 0.27 mmol) in DMF (1.4 mL) at room temperature. The mixture was stirred at 90 °C for 1.5 h then was cooled to room temperature and concentrated under reduced pressure. The residue was purified by flash chromatography (hexane:EtOAc, 8:2 to 7:3) to give the title compound (101 mg, 99% yield).

**g) 5,11-Dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one**

A stream of ozonised oxygen was bubbled through a cold (-78 °C) solution of 5,11-dihydro-11-ethyl-5-methyl-8-(2-propenyl)-2-(trifluoromethyl)-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one (100 mg, 0.27 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.7 mL) and MeOH (2.7 mL) for 2.5 h. A stream of N<sub>2</sub> was next bubbled through the solution for 15 min and then solid NaBH<sub>4</sub> (52 mg, 1.4 mmol) was added to the solution. The reaction mixture was allowed to warm to room temperature. After 1 h, aqueous saturated NH<sub>4</sub>Cl (5 mL) was added and the mixture was stirred at room temperature for 2 h. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> and the aqueous layer was re-extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was successively washed with water and brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:hexane: 75:25 to EtOAc 100%) to give the title compound (60 mg 59% yield) as a white solid.

**h) 5,11-Dihydro-11-ethyl-5-methyl-8-[2-(4-quinolinylloxy)ethyl]-2-(trifluoromethyl)-6H-dipyrido[3,2b:2',3'-e] [1,4]diazepin-6-one**

Diethyl azodicarboxylate (DEAD) (38  $\mu$ L, 0.24 mmol) was added dropwise to

- a solution of 5,11-dihydro-11-ethyl-8-(2-hydroxyethyl)-5-methyl-2-(trifluoromethyl)-6*H*-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one (58 mg, 0.16 mmol), 4-hydroxyquinoline (35 mg, 0.24 mmol) and Ph<sub>3</sub>P (62 g, 0.24 mmol) in THF (1.6 mL) at room temperature. The mixture was stirred at room
- 5 temperature for 1h then was concentrated under reduced pressure. The residue was purified by flash chromatography (EtOAc:MeOH; 95:5) to give the title compound (24 mg, 31% yield) as a white solid: MS (ESI) *m/z* 494, 495 (MH)<sup>+</sup>.
- 10 i) **5,11-Dihydro-11-ethyl-5-methyl-8-[2-[(1-oxido-4-quinolinyl)oxy]ethyl]-2-(trifluoromethyl)-6*H*-dipyrido[3,2-*b*:2',3'-*e*] [1,4]diazepin-6-one**  
Solid mCPBA (80-85%, *m*-chloroperbenzoic acid) (16 mg, 0.08 mmol) was added to a solution of 5,11-dihydro-11-ethyl-5-methyl-8-[2-(4-quinolinylloxy)ethyl]-2-(trifluoromethyl)-6*H*-dipyrido[3,2-*b*:2',3'-*e*]
- 15 [1,4]diazepin-6-one (22 mg, 0.04 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL) at room temperature. The mixture was stirred at room temperature for 2.5 h then diluted with CH<sub>2</sub>Cl<sub>2</sub>. The resulting solution was successively washed with aqueous 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, aqueous saturated NaHCO<sub>3</sub> and brine, dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The residue was
- 20 purified by flash chromatography (MeOH:CHCl<sub>3</sub>, 5% to 10%) to give the title compound (11 mg, 52% yield) as a white solid: MS (ESI) *m/z* 511 (MH)<sup>+</sup>.

51

**TABLE 1**  
Compounds of formula I




Table 1 Entry #	R <sup>2</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>11</sup>	Q
1	Cl	Me	H	Et	
2	Cl	H	Me	Et	
3	H	Me	H	CycPr	
4	F	H	Me	Et	
5	F	Me	H	CycPr	
6	Cl	H	Me	CycPr	
7	F	Me	H	Et	

52





Table 1 Entry #	R <sup>2</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>11</sup>	Q
8	H	Me	H	Et	
9	F	Me	H	Et	
10	Cl	H	Me	CycPr	
11	F	H	Me	Et	
12	H	Me	H	Et	
14	Cl	Me	H	Et	
15	H	Me	H	CycPr	

53



I.

Table 1 Entry #	R <sup>2</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>11</sup>	Q
17	Cl	H	Me	Et	
18	F	H	Me	Et	
19	Cl	Me	H	Et	
20	Cl	H	Me	Et	
21	H	H	Me	Et	
22	Cl	H	H	Et	
23	F	Me	H	Et	



54




Table 1 Entry #	R <sup>2</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>11</sup>	Q
24	H	H	Me	Et	
25	Cl	Me	H	Et	
26	H	H	Et	Et	
27	H	H	Et	Et	
28	Me	H	Me	Et	
29	CF <sub>3</sub>	H	H	Et	

## REVERSE TRANSCRIPTASE (RT) ASSAYS

### ASSAY THEORY:

- 5 Among the enzymes for which Human Immunodeficiency Virus (HIV-1) encodes is a reverse transcriptase (1), so-named because it transcribes a DNA copy from an RNA template. This activity can be quantitatively measured in a cell-free enzyme assay and is based upon the observation that reverse transcriptase is able to use a synthetic template poly r(C) primed with a
- 10 biotinylated oligo d(G) to transcribe a radio-labelled DNA strand utilising 3H-dGTP as a substrate. The assay described below utilises the wild-type enzyme (which is the predominant form of the enzyme observed in patients infected with HIV-1) and can also be used with mutant RT enzymes (for example, Y181C, prepared by site-directed mutagenesis in which the tyrosine
- 15 residue at codon 181 has been replaced by a cysteine residue) in analogous assay conditions. This assay allows compounds to be evaluated for their effectiveness at inhibiting the mutant enzymes.

### MATERIALS:

20

#### a) Preparation of the enzyme

- Some HIV-1 IIIB clone BH10 RT mutants were provided by Dr. C.-K. Shih (Boehringer Ingelheim Pharmaceuticals Inc., U.S.A.) in the vector pKK233-2 (Pharmacia). Briefly an HIV RT clone pKRT2 containing only the RT p66
- 25 gene regulated by the lac operon/trc promoter was obtained from Dr. W. Summers (Yale University) (2). A variety of specific amino acid substitutions were introduced into the wild-type RT gene by site-directed mutagenesis. RT clones were subcloned into the pKK233-2 bacterial expression vector. Clones provided included wild-type, Val106Ala, Tyr181Cys, Tyr188Cys,

Tyr188Leu, Gly190Ala and Pro236Leu. Others were made in-house by site-directed mutagenesis of the pKK233-2 RT clones including Lys103Asn, Lys103Asn/Tyr181Cys, Lys103Asn/Leu100Ile, Lys103Asn/Pro225His, and Lys103Asn/Val108Ile.

5

#### b) Purification of Enzyme

Purification of recombinant reverse transcriptase was performed using a combination of methods previously described (3). A single colony from a fresh plate of transformed JM109 cells was used to initiate growth of a pre-culture grown o/n at 37°C. Two liters of growth medium were inoculated with this pre-culture. At OD<sub>600</sub> ~ 1.5 (5-6 h at 37°C), RT gene expression was induced with IPTG (1 mM final), and the fermentation was continued for a few more hours at 37 °C. After centrifugation, supernatants were discarded while cell pellets were pooled and stored at -80 °C until purification. Cells were thawed at 4 °C overnight and suspended in lysis buffer (MES 50mM pH 6, EDTA 1mM, 10% v/v glycerol, 0.02% w/v OBG, 0.02% w/v sodium azide). Lysozyme was added and the mixture was incubated on ice for 40 minutes. After homogenization using a Dounce in presence of lysozyme and sonication, the cells were centrifuged for 30 minutes. Supernatant (S1) was saved and stored at 4°C. The centrifuged pellet was resuspended in extraction buffer (MES 50mM pH 6, KPO<sub>4</sub> 50 mM pH 6, KCl 100mM, 10% v/v glycerol, 0.02% w/v OBG, 0.02% w/v sodium azide) and stirred for 30 minutes at 4°C. This second mixture was centrifuged again and the supernatant (S2) was saved. The above procedure was repeated 2 more times saving supernatants S3 and S4 and one last extraction was performed overnight (S5). Polymyxin B (0.005% final) was added to the combined supernatants to remove nucleic acids. This solution was stirred for 75 minutes at 4 °C and centrifuged for 1 h. The supernatant (SS1) was

10

15

20

25

precipitated on ice with 20% w/v ammonium sulfate and stirred for 1 h at 4 °C. The mixture was then centrifuged and the resulting supernatant (SS2) was precipitated with additional 40% w/v ammonium sulfate (60% total), stirred for 1 h and centrifuged again. The final pellet (P1) was stored  
5 overnight at 4 °C before undergoing purification the following day. All steps of the purification were performed at 4 °C unless otherwise stated.

Pellet (P1) was resuspended into MES 50mM pH 6, KPO<sub>4</sub> 10 mM pH 6, KCl 100mM, 10% v/v glycerol, 0.02% w/v OBG, 0.02% w/v sodium azide. The suspension was dialysed against the same buffer overnight using 12-14kD  
10 MWCO dialysis tubing. The dialysate was centrifuged and the supernatant was filtered through Millex-PF 0.8 µm filter units. The filtered sample was loaded on a Hydroxy Apatite column (30-mL bed volume) and washed with the same buffer. The bound enzyme was eluted with ~220 mL of a linear gradient of 10 to 300mM KPO<sub>4</sub> in the above buffer. The fractions containing  
15 p66/p51 heterodimer (as determined by SDS-PAGE 8% and Western blotting) were pooled for the next column. The RT containing fractions were diluted two-fold with Bis-Tris propane 50 mM pH 7.0, 0.02% w/v OBG, 10% v/v glycerol, 0.02% w/v sodium azide and loaded on a Hi-Trap Heparin Sepharose column (5-mL bed volume) and washed with the same buffer. The  
20 bound RT was then eluted with 75 mL of a linear gradient of 0 to 1 M ammonium sulfate in the same buffer. RT-containing fractions were pooled according to SDS-PAGE and Western blotting analyses. Protein concentration of this pool was determined by the Bradford method using BSA as standard. The final enzyme preparation was dialyzed in MES 50mM  
25 pH 6, KPO<sub>4</sub> 300 mM pH 6, KCl 175mM, 10% v/v glycerol, 0.02% w/v sodium azide and aliquoted and frozen at -80° C.

**ASSAY PROCEDURE:**

The radiometric enzyme assay has been adapted to a 96-well microtiter plate format and uses streptavidin scintillation proximity beads. The assay is briefly described below. The HIV-1 RT enzyme was thawed and appropriately diluted into Tris/HCl 50 mM pH 7.8 containing NaCl 60mM, MgCl<sub>2</sub> hexahydrate 2mM, DTT 6mM, GSH 2mM and 0.02% w/v Chaps to give  $\approx 3$  nM enzyme. To 30  $\mu$ L of this enzyme solution was added 10  $\mu$ L of inhibitor solution (50  $\mu$ M to 2.5 nM inhibitor in same assay buffer as above containing 15% v/v DMSO). The plate was pre-incubated for 15 minutes at room temperature before proceeding to the next step. In this pre-incubation step, the highest and lowest inhibitor concentrations were 12.5  $\mu$ M and 0.62 nM respectively and the concentration of DMSO was 3.75% v/v. Then the enzymatic reaction was initiated by addition of 10  $\mu$ L of substrates solution. The final reaction mixture contained Tris/HCl 50 mM pH 7.8, NaCl 60 mM, MgCl<sub>2</sub>•6H<sub>2</sub>O 2 mM, DTT 6 mM, GSH 2 mM, Chaps 0.02% w/v DMSO 3% v/v, Poly rC 179 nM, Biotin dG<sub>15</sub> 18 nM, dGTP 288 nM, <sup>3</sup>H-dGTP 71nM, and 1-2 nM enzyme.

In this incubation step, the highest and lowest inhibitor concentrations were 10  $\mu$ M and 0.5 nM respectively. After addition of substrates, the plate was covered with a plastic seal and incubated for 1 hour at 37°C in a dry incubator. Then the reaction was quenched by addition of 75  $\mu$ L of EDTA 0.5M containing 5mg/mL of streptavidin scintillation proximity beads.

The plate was shaken for 2 minutes at medium speed and incubated 1 hour at room temperature. Then 75  $\mu$ L of cesium chloride 7 M solution was added, the plate was shaken for 2 minutes at medium speed and incubated again for 1 hour at room temperature. The plate was then covered with a plastic seal and counted using the TopCount-NXT™ Microplate Scintillation & Luminescence Counter, (Packard). Each well was counted for 60 seconds.

Each row contained at its extremities a blank and a control well.

The calculation for percent inhibition is as follows:

$$\% \cdot inhibition = \left( 1 - \left[ \frac{cpm \cdot well - cpm \cdot blank}{cpm \cdot control - cpm \cdot blank} \right] \right) * 100$$

Using the above assay, compounds of the invention were tested for  
5 inhibition of RT wild-type (WT) and mutant enzymes. The results are listed  
in Table 2, as IC<sub>50</sub> (nM).

To confirm the ability of these compounds to inhibit HIV replication, they  
were also tested in the human T-Cell Culture (Syncytia) Assay described  
10 below.

#### **ELISA assay for assessment of activity in cell culture**

Compounds of the invention were tested for their ability to inhibit HIV  
15 replication in cell culture in a 96-well plate assay. Complete RPMI 1640,  
consisting of RPMI 1640 + 10% fetal bovine serum, 10 µg/ml gentamycin and  
10 µM β-mercaptoethanol was used for dilution of the compound as well as  
cell growth medium. The T lymphocyte cell line C8166 was infected at a  
multiplicity of infection of 0.001 with viruses coding for wild-type and  
20 mutant reverse transcriptase. Cells were then incubated for three days in the  
presence of serial dilutions of the compounds of the invention. The  
supernatant was pooled from eight replica wells and the concentration of  
extracellular p24 was determined using a commercially available HIV-1 p24  
antigen assay kit (Beckman-Coulter®). The level of inhibition (% inhibition)  
25 was calculated with the following equation:

$$\% \cdot inhibition = \left( 1 - \left[ \frac{p24 pg / mL \cdot inhibitor}{p24 pg / mL \cdot control} \right] \right) * 100$$

The results are listed in Table 3, as EC<sub>50</sub> (nM).

### Specificity Assays

5

In order to assess the specificity of the enzyme inhibitory activity of the compounds provided by the invention, a few were tested, for their ability to inhibit other viral polymerases (such as Hepatitis C virus and Respiratory Syncytial virus RNA-dependent RNA polymerases) as well as mammalian  
10 polymerases (such as Calf Thymus RNA-dependent DNA polymerase and Human telomerase) using known assay methods. None of the compounds so tested was observed to possess any significant inhibitory activity against these enzymes. These results indicate that the enzyme inhibitory activity of the compounds provided by the invention is directed rather specifically  
15 against HIV-1 RT.

### REFERENCES (INCORPORATED HEREIN BY REFERENCE)

1. Benn, S., et al. Science 230:949, 1985.
2. D'Aquila, R. T. and Summers, W. C. J. Acq. Imm. Def. Syn. 2:579, 1989.
- 20 3. a) Warren, T. C. et al. Protein Expression & Purification 3:479, 1992; b) Kohlstaedt, L. A. Science 256(5065): 1783, 1992.

TABLE 2  
Inhibition of Wild-type and mutant strains of RT for compounds of formula I

compound no. (see Table 1)	IC <sub>50</sub> WT RT (nM)	IC <sub>50</sub> K103N (nM)	IC <sub>50</sub> Y181C (nM)	IC <sub>50</sub> V106A (nM)	IC <sub>50</sub> P236L (nM)	IC <sub>50</sub> Y188L (nM)	IC <sub>50</sub> G190A (nM)	IC <sub>50</sub> K103N/ Y181C (nM)	IC <sub>50</sub> K103N/ P225H (nM)	IC <sub>50</sub> Y188C (nM)	IC <sub>50</sub> K103N/ V108I (nM)	IC <sub>50</sub> K103N/ L100I (nM)
1	10.5	19	20	132		2740		52	31.2		29	28.5
2	5.5	28.5	25	148	50.5	258	14.5	24.5	27	7.1	31	46
3	16.5	45	73	79		3236		612	56		47	124
4	11	22.5	26	206		520		52	27		32	108
5	19	37.5	76	61		2541		536	38		38	98
6	25											
7	9.8	26	38	74		4318		199	46		45	98
8	10	37	43	190		7511		301	67		63	135
9	14	23	43	98		3395		175	25		58	88
10	41	67	206	1372		1422		710			255	229
11	9.5	18	27	336	56	661	19	55	15	6.6	55	102
12	12	31	38	121		>10000		212	38		75	124
14	15	19	28	77		2298		61	30		59	24
15	61	66	226	206		>10000		1522	93		215	270
17	14	25	29	319	60	198	25	48	23	8.1	83	40
18	11	37	41	737		1167		146	20		96	158
19	7	25	20	89		3938		91	16		56	27



compound no. (see Table 1)	IC <sub>50</sub> WT RT (nM)	IC <sub>50</sub> K103N (nM)	IC <sub>50</sub> Y181C (nM)	IC <sub>50</sub> V106A (nM)	IC <sub>50</sub> P236L (nM)	IC <sub>50</sub> Y188L (nM)	IC <sub>50</sub> G190A (nM)	IC <sub>50</sub> K103N/ Y181C (nM)	IC <sub>50</sub> K103N/ P225H (nM)	IC <sub>50</sub> Y188C (nM)	IC <sub>50</sub> K103N/ V108I (nM)	IC <sub>50</sub> K103N/ L100I (nM)
20	10	36	22	661		439		51	17		64	44
21	17	44	51	1106	129	1357	26	95	40	9.9	120	168
22	45			2493		2990		336				
23	6.1			105		5614		247				
24	5.6			344		627	25	70				169
25	8	19	17	84		2537		64	23		38	28
26	8.9	40	51	1745		5204		200	85		147	443
27	38			2271		5335		299				
28	19			605		1190	25	141	25	9.0	35	28
29	35			893		578		84				

**TABLE 3**  
**Inhibition of wild type and mutant strains of HIV in cell cultures by compounds of formula I**


compound no. (see Table 1)	EC <sub>50</sub> WT RT (nM)	EC <sub>50</sub> K103N (nM)	EC <sub>50</sub> Y181C (nM)	EC <sub>50</sub> V106A (nM)	EC <sub>50</sub> Y188L (nM)	EC <sub>50</sub> K103N/ Y181C (nM)	EC <sub>50</sub> K103N/ P225H (nM)	EC <sub>50</sub> K103N/ V108I (nM)	EC <sub>50</sub> K103N/ L100I (nM)
1	2.6								
2	2.2								
3	1.3								
4	0.98								
5	0.83								
6									
7	1.1								
8	0.93								
9	0.92			5	118	7.6			
10									
11	0.34	2	4.1	28	52	7.5	2.0	3.9	3.2
12	0.26	2.8	2.7	12.4			1.5	5.0	2.9
14	0.67								
15									
17	0.34	1.5	2.8	24	58	2.3	1.2	2.7	1.2
18	0.34					9.5			
19	0.37					22			

compound no. (see Table 1)	EC <sub>50</sub> WT RT (nM)	EC <sub>50</sub> K103N (nM)	EC <sub>50</sub> Y181C (nM)	EC <sub>50</sub> V106A (nM)	EC <sub>50</sub> Y188L (nM)	EC <sub>50</sub> K103N/ Y181C (nM)	EC <sub>50</sub> K103N/ P225H (nM)	EC <sub>50</sub> K103N/ V108I (nM)	EC <sub>50</sub> K103N/ L100I (nM)
20	0.40					11			
21	0.53	3.5	4.8	65	67	4.3	2.9	5.0	5.9
22									
23									
24									
25	2.2								
26	2.2								
28	1.3	2.3	2.6	42	70	7.5			
29	1.6			83	65	3.5			

65

**WHAT IS CLAIMED IS:**

1. A compound of the general formula I:



- 5 wherein

R² is selected from the group consisting of H, F, Cl, C<sub>1-4</sub> alkyl, C<sub>3-4</sub> cycloalkyl and CF<sub>3</sub>;


R⁴ is H or Me;

R⁵ is H, Me or Et, with the proviso that R⁴ and R⁵ are not both Me, and if R⁴ is

- 10 Me then R⁵ cannot be Et;

R¹¹ is Me, Et, cyclopropyl, propyl, isopropyl, or cyclobutyl; and

Q is selected from the group consisting of:




or a pharmaceutically acceptable salt thereof.

66


2. A compound according to claim 1, wherein R<sup>11</sup> is Et or cyclopropyl or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 1, wherein Q is selected from the  
5 group consisting of:



or a pharmaceutically acceptable salt thereof.

4. A compound according to claim 1, wherein Q is:



10

or a pharmaceutically acceptable salt thereof.


5. A compound according to claim 1, wherein R<sup>5</sup> is Me,  
or a pharmaceutically acceptable salt thereof.

15

6. A compound according to claim 1, wherein R<sup>11</sup> is Et or a pharmaceutically acceptable salt thereof.


7. The compound:

67




or a pharmaceutically acceptable salt thereof.

8. The compound:



5 or a pharmaceutically acceptable salt thereof.


9. The compound:



or a pharmaceutically acceptable salt thereof.

10. An inhibitor of HIV replication of the general formula I:

68



I

wherein


R<sup>2</sup> is selected from the group consisting of H, F, Cl, C<sub>1-4</sub> alkyl, C<sub>3-4</sub> cycloalkyl and CF<sub>3</sub>;

5 R<sup>4</sup> is H or Me;

R<sup>5</sup> is H, Me or Et, with the proviso that R<sup>4</sup> and R<sup>5</sup> are not both Me, and if R<sup>4</sup> is Me then R<sup>5</sup> cannot be Et;

R<sup>11</sup> is Me, Et, cyclopropyl, propyl, isopropyl, or cyclobutyl;; and

Q is selected from the group consisting of:




10

or a pharmaceutically acceptable salt thereof.


11. An inhibitor of HIV replication according to claim 10, wherein R<sup>11</sup> is Et or cyclopropyl or a pharmaceutically acceptable salt thereof.

12. An inhibitor of HIV replication, according to claim 10, wherein Q is selected from the group consisting of:



- 5 or a pharmaceutically acceptable salt thereof.

- 13.. An inhibitor of HIV replication, according to claim 10, wherein Q is:



- or a pharmaceutically acceptable salt thereof.

10


14. An inhibitor of HIV replication, according to claim 10, wherein R<sup>5</sup> is Me.

- 15 15. An inhibitor of HIV replication, according to claim 10, wherein R<sup>11</sup> is Et.

16. An inhibitor of HIV replication of the formula:




70




or a pharmaceutically acceptable salt thereof.

17. An inhibitor of HIV replication of the formula:



or a pharmaceutically acceptable salt thereof.

18. An inhibitor of HIV replication of the formula:



or a pharmaceutically acceptable salt thereof.

19. A method for the treatment or prevention of HIV infection, comprising administering to a patient an HIV inhibiting amount of a

compound of formula I according to claim 1, or a pharmaceutically acceptable salt thereof.

20. A pharmaceutical composition for the treatment or prevention of HIV  
5 infection, comprising a compound of formula I according to claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/CA 01/00890

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D471/14 A61K31/55 //(C07D471/14,243:00,221:00,221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, PAJ, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 767 172 A (BOEHRINGER INGELHEIM PHARMA) 9 April 1997 (1997-04-09) examples 79,85,96,97 & US 5 705 499 A 6 January 1998 (1998-01-06) cited in the application	1-20
Y	KLUNDER, JANICE M. ET AL: "Novel Nucleoside Inhibitors of HIV-1 Reverse Transcriptase. 7. 8-Arylethylidipyridodiazepinones as Potent Broad-Spectrum Inhibitors of Wild-Type and Mutant Enzymes" J. MED. CHEM. (1998), 41(16), 2960-2971 , XP002181339 cited in the application compounds 10,18 - table 1; 41,43 -table 3 --- -/-	1-20

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

30 October 2001

Date of mailing of the international search report

13/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Frelon, D

## INTERNATIONAL SEARCH REPORT

International Application No

PCI/CA 01/00890

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>CYWIN, CHARLES L. ET AL: "Novel Nucleoside Inhibitors of HIV-1 Reverse Transcriptase. 8. 8-Aryloxymethyl- and 8-Arylthiomethyldipyridodiazepinones" J. MED. CHEM. ( 1998 ), 41(16), 2972 -2984</p> <p>' XP002181340 compounds II, 27-29, 32-34, 43 - table 1 -----</p>	1-20

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 01/00890

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0767172	A	09-04-1997	CA 2187146 A1	07-04-1997
			EP 0767172 A1	09-04-1997
			JP 9188680 A	22-07-1997
			US 5705499 A	06-01-1998
<hr/>				